

# Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.


Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

5. As there are  $2^{2}(n-1)(n-2)$  different arcs on  $c_{0}$ , the total number x of polygons is

$$2^{x}(n-1)(n-2)(n-3)!2^{2n-3},$$
  
or  $x=(n-1)!2^{2n-1}.$ 

6. To find the total number of all possible spherical polygons that may be formed by arcs of n great circles, it is clear that there will be  $(r-1)!2^{2r-1}$  polygons of r sides, and consequently



$$\sum_{r=3}^{r=n} {n \choose r} (r-1)! 2^{2r-1}$$

will be the number of all possible polygons.

7. Examples. With three great circles on a sphere  $(3-1)!2^{6-1}=64$  triangles may be formed. The number of spherical triangles and quadrangles which may be formed by four great circles is

$$(4-1)!2^{8-1}+{}^{4}C_{3}.(3-1)!2^{6-1}=768+4.64=1024.$$

## PROBLEMS FOR SOLUTION.

#### ALGEBRA.

226. Proposed by ELMER SCHUYLER, Brooklyn, N. Y.

Find the real roots of the system

$$x^{2}+w^{2}+v^{2}=a^{2}$$
,  $vw+u(y+z)=bc$ ,  
 $w^{2}+y^{2}+u^{2}=b^{2}$ ,  $wu+v(z+x)=ca$ ,  
 $v^{2}+u^{2}+z^{2}=c^{2}$ ,  $uv+w(x+y)=ab$ .

227. Proposed by G. I. HOPKINS, A. M., Manchester, N. H.

Solve  $x + y + xy + x^2y + xy^2 + x^3y + 2x^2y^2 + xy^3 + x^3y^2 + x^2y^3 = 11$ ;  $x^4y + 3x^3y^2 + 3x^2y^3 + 2x^4y^2 + 4x^2y^3 + 2x^2y^4 + 4x^4y^3 + 4x^3y^4 + xy^4 + x^5y^2 + x^5y^3 + 2x^4y^4 + x^2y^5 + x^3y^5 = 30$ .

#### GEOMETRY.

### 251. Proposed by R. D. CARMICHAEL, Hartselle, Ala.

Represent the vertices of any regular polygon by the consecutive numbers 1, 2....p....q....r.. To find the sides and area of the triangle formed by joining p, q, and r.